

AD-A169 896 NOTE ON A CHARACTERIZATION OF EXPONENTIAL DISTRIBUTIONS 1/1  
(U) MARYLAND UNIV COLLEGE PARK DEPT OF MANAGENET  
SCIENCES AND STATISTICS S KOTA ET AL. JUL 86  
UNCLASSIFIED UM/HSS-86/4 N00014-84-K-0301 F/G 12/1 NL





1.0



1.1



1.25



1.4

2.8



2.5

3.15



2.2

3.5



2.0

4.0



1.8

4.5



1.6

## Note on a characterization of exponential distributions

S. Kotz,  
University of Maryland \*)  
F.W. Steutel,  
Eindhoven University of Technology \*\*)

JULY 1986

DTIC  
ELECTE  
JUL 17 1986  
S D E

## ABSTRACT

Let  $U$  be uniformly distributed on  $(0,1)$  and let  $Y$  and  $Y' \stackrel{d}{=} Y$  be random vectors with nonnegative components,  $U, Y$  and  $Y'$  independent. It is shown that the relation  $Y \stackrel{d}{=} U(Y+Y')$  is satisfied if and only if the components of  $Y$  are multiples of a single exponentially distributed random variable.

## 1. One-dimensional case

In the solution to problem 159 in [3] the following question is answered. Let  $U, Y^{(1)}$  and  $Y^{(2)}$  be independent random variables,  $U$  uniformly distributed on  $(0,1)$ ,  $Y^{(1)}$  and  $Y^{(2)}$  distributed as  $Y$ . For what distributions of  $Y$  is it true that

$$(1) \quad Y \stackrel{d}{=} U(Y^{(1)} + Y^{(2)}) \quad ?$$

There is a two-parameter family of solutions (cf. [3]), but under the additional assumption that  $Y$  is nonnegative, (1) characterizes the *exponential distributions*. We state this result as a proposition, and give a proof along the lines of the proof in [3].

**Proposition 1.** Let  $Y \geq 0$  with probability 1 and let  $Y$  satisfy condition (1). Then  $Y$  has an exponential distribution (possibly concentrated at zero).

**Proof** If  $\phi$  denotes the Laplace-Stieltjes transform (LST) of the distribution of  $Y$ , i.e.  $\phi(s) = E \exp(-sY)$ , then (1) is equivalent to

$$(2) \quad \phi(s) = \int_0^1 \phi(us) du = \frac{1}{s} \int_0^s \phi^2(t) dt.$$

Since  $\phi$  and  $\phi^2$  are LST's, they are differentiable for  $s > 0$ . Differentiation of (2) yields

\*) Postal address: College of Business and Management, University of Maryland, College Park, Maryland 20742.

\*\*) Postal address: Department of Mathematics and Computing Science, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<i>per</i>
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

$$(3) \quad s\phi'(s) + \phi(s) = \phi^2(s).$$

Substitution of  $\phi = (f+1)^{-1}$  leads to

$$f'(s)/f(s) = 1/s$$

with solution  $f(s) = as$  and so

$$(4) \quad \phi(s) = \frac{1}{1+as}.$$

where  $a \geq 0$ . This proves the proposition □

In exactly the same way the following generalization can be proved.

**Proposition 2.** If  $U, Y^{(1)}, \dots, Y^{(N)}$  are independent,  $U$  uniformly distributed on  $(0,1)$  and the  $Y^{(i)}$  distributed as  $Y$ , then

$$(5) \quad Y \stackrel{d}{=} U(Y^{(1)} + \dots + Y^{(N+1)})$$

if and only if the LST  $\phi_N$  of  $Y$  is of the form

$$(6) \quad \phi_N(s) = \frac{1}{1+as^{1/N}},$$

where  $a \geq 0$ .

**Remark.** Since  $\exp(-s^{1/N})$  is an infinitely divisible (even stable; see [1], p. 448) LST it follows from Theorem 2 in [4] that  $\phi_N$  in (6) is indeed the LST of a (infinitely divisible) probability distribution having no moments, of course. One can even make  $N$  a continuous variable:  $S(1) \stackrel{d}{=} U S(t+1)$ , where  $S(\cdot)$  is a process with nonnegative stationary and independent increments. Then  $S(1)$  must have an LST of the form  $(1+as^{1/t})^{-1}$  with  $t > 0$ .

An other generalization is considered in the next section.

## 2 Multi-dimensional case

Now let  $Y$  be an  $n$ -dimensional random vector with nonnegative components:

$$Y = (Y_1, \dots, Y_n).$$

and as before let  $U, Y^{(1)}$  and  $Y^{(2)}$  be independent with  $U$  uniform on  $(0,1)$  and  $Y^{(1)}$  and  $Y^{(2)}$  distributed as  $Y$ . Now let

$$Y \stackrel{d}{=} U(Y^{(1)} + Y^{(2)}).$$

where addition is component-wise, and let the  $n$ -dimensional LST  $\phi$  be defined by

$$(7) \quad \phi(s_1, \dots, s_n) = E \exp \left[ - \sum_{j=1}^n s_j Y_j \right].$$

Then in exactly the same way as in (2) we have

$$\phi(s_1, \dots, s_n) = \int_0^1 \phi^2(us_1, \dots, us_n) du.$$

or putting  $s_j = \alpha_j s$ .

$$s \phi(\alpha_1 s, \dots, \alpha_n s) = \int_0^s \phi^2(\alpha_1 t, \dots, \alpha_n t) dt.$$

Writing  $\phi(\alpha_1 s, \dots, \alpha_n s) = \phi_\alpha(s)$  for all  $\alpha \in \mathbb{R}_+^n$  we obtain

$$s \phi_\alpha(s) = \int_0^s \phi_\alpha^2(t) dt.$$

the same equation as (2). It follows that for all  $\alpha \in \mathbb{R}_+^n$  We have

$$(8) \quad \phi_\alpha(s) = \phi(\alpha_1 s, \dots, \alpha_n s) = (1 + a(\alpha)s)^{-1}.$$

where  $a(\alpha) = a(\alpha_1, \dots, \alpha_n)$  by the definition of  $\phi_\alpha$  satisfies

$$(9) \quad a(s\alpha_1, \dots, s\alpha_n) = s a(\alpha_1, \dots, \alpha_n).$$

i.e.  $a(\alpha)$  is homogeneous of degree one. We are now ready to prove

Proposition 3. A random vector  $Y = (Y_1, \dots, Y_n)$  with  $Y^j \geq 0$  ( $j = 1, \dots, n$ ) satisfies

$$(10) \quad Y \stackrel{d}{=} U(Y^{(1)} + Y^{(2)})$$

with  $U, Y^{(1)}$  and  $Y^{(2)}$  independent,  $U$  uniform on  $(0,1)$  and  $Y^{(1)} \stackrel{d}{=} Y^{(2)} \stackrel{d}{=} Y$  if and only if the LST  $\phi$  of  $Y$  is of the form

$$(11) \quad \phi(s_1, \dots, s_n) = \frac{1}{1 + a_1 s_1 + \dots + a_n s_n}.$$

where  $a_1 \geq 0, \dots, a_n \geq 0$ .

Proof. From (7) and (8) with  $s_j = \alpha_j s$  and (9) it follows that for all  $(\alpha_1, \dots, \alpha_n) \in \mathbb{R}_+^n$

$$(12) \quad \alpha_1 Y_1 + \dots + \alpha_n Y_n \stackrel{d}{=} a(\alpha_1, \dots, \alpha_n) X.$$

where  $X$  is exponentially distributed with expectation one and  $Y_1, \dots, Y_n$  are exponential with expectations  $a(1,0, \dots, 0), \dots, a(0, \dots, 0,1)$ . Taking expectations in (11) we obtain

$$(13) \quad \alpha_1 a(1, 0, \dots, 0) + \dots + \alpha_n a(0, \dots, 0, 1) = a(\alpha_1, \dots, \alpha_n).$$

If we put  $a(1, 0, \dots, 0) = a_1, \dots, a(0, \dots, 0, 1) = a_n$ , then (11) follows from (8) and (13).

Remark 1. From proposition 3 it follows that the only random vectors  $Y = (Y_1, \dots, Y_n)$  satisfying (10) are of the form

$$Y = (a_1 X, \dots, a_n X),$$

where  $X$  is an exponentially distributed random variable with expectation one. This means that  $Y$  has a (singular) exponential distribution concentrated on the ray with direction  $(a_1, \dots, a_n)$  through the origin. So, none of the classical multivariate distributions, such as described in [2] satisfy (10).

Remark 2. One could also generalize (5) to  $n$ -dimensional vectors; this leads to results similar to proposition 3.

Remark 3. If the condition  $Y_1 \geq 0, \dots, Y_n \geq 0$  is dropped then more general solutions than (11) are possible. For  $n = 2$ , for instance, (10) is satisfied for  $Y$  with a characteristic function of the form

$$\Psi(t_1, t_2) = (1 + a_0 \sqrt{t_1^2 + t_2^2} - a_1 i t_1 - a_2 i t_2)^{-1},$$

with  $a_0 \geq 0, a_1$  and  $a_2$  real. A similar situation occurs for  $n = 1$  (cf. [3]).

### References

- [1] Feller, W. (1971) *An introduction probability theory and its applications*, vol. 2. Wiley, New York, etc.
- [2] Johnson, N.L. and Kotz, S. (1972) *Distributions in Statistics: Continuous Multivariate Distributions*, Wiley, New York.
- [3] Problem Section (1985). *Statistica Neerlandica* 39, 57-58.
- [4] Steutel, F.W. (1968) A class of infinitely divisible mixtures. *Ann. Math. Statist.* 39, 1153-1158.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT		
2b DECLASSIFICATION / DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) <b>UMHSS/86/4</b>			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION <b>University of Maryland</b>		6b OFFICE SYMBOL (If applicable)	7a NAME OF MONITORING ORGANIZATION <b>Office of Naval Research</b>		
6c ADDRESS (City, State, and ZIP Code) <b>Dep't. of Management and Statistics University of Maryland College Park, Md. 20742</b>			7b ADDRESS (City, State, and ZIP Code)		
8a NAME OF FUNDING / SPONSORING ORGANIZATION <b>CNR</b>		8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <b>NCOC14-84-K-0301</b>		
8c ADDRESS (City, State, and ZIP Code) <b>Stat. and Probability Program (CODE 436) Office of Naval Research Arlington, Va. 22217</b>			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO	PROJECT NO	TASK NO
			WORK UNIT ACCESSION NO		
11 TITLE (Include Security Classification) <b>Note on a characterization of exponential distributions</b>					
12 PERSONAL NAME(S) <b>Folz, Samuel and Stentel, R.W.</b>					
13a TYPE OF REPORT <b>technical</b>		13b TIME COVERED FROM <b>9/85</b> TO <b>8/86</b>		14 DATE OF REPORT (Year, Month, Day) <b>July 1986</b>	
15 PAGE COUNT <b>4</b>					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	<b>exponential distribution; uniform distribution; Laplace-Stieltjes transform; infinite divisibility</b>		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <b>Let <math>U</math> be uniformly distributed on <math>(0,1)</math> and let <math>Y</math> and <math>Y' \stackrel{d}{=} Y</math> be random vectors with nonnegative components, <math>U</math>, <math>Y</math> and <math>Y'</math> independent. It is shown that the relation <math>Y \stackrel{d}{=} U(Y+Y')</math> is satisfied if and only if the components of <math>Y</math> are multiples of a single exponentially distributed random variable.</b>					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		
22a NAME OF RESPONSIBLE INDIVIDUAL <b>Samuel Folz</b>			22b TELEPHONE (Include Area Code) <b>301 454-6805</b>		22c OFFICE SYMBOL

END

DT/C

8-86